

Growth Response Of Brassica Oleracea Var. Botrytis L. To Carabao Manure Boosted With Carabao Manure Tea As Bio-Organic Fertilizer

Leah C. Tuan ¹

¹Department of Agricultural Sciences,
College of Agriculture, Fisheries, and Natural Resources,
University of Eastern Philippines, University Town,
Northern Samar, Philippines
leahtuanc@gmail.com

Abstract

Utilization of carabao manure as a sustainable approach on nutrient management for cauliflower production was studied for three months between the months of October 2016 to January 2017 at the University Demo Farm, University of Eastern Philippines, Main Campus, University Town, Catarman, Northern Samar, Philippines. The main objective was to determine the effect of carabao manure on the growth of cauliflower. Carabao manure preparations included dried carabao manure, administered at 1.0 kg and 1.5 kg levels boosted with manure tea. Growth performance of cauliflowers were significantly influenced by dried carabao manure applications boosted with manure tea application. Better performance was observed in plants treated with 1.5 kg dried carabao manure than with 1.0 kg was evident on the parameters evaluated, indicative of the nutrient sufficiency of the bio-fertilizer. Carabao manure, a farm waste product, an untapped resource, has been proven to be an effective, safe and an economical bio-organic fertilizer. A commodity capable of becoming a promising industry; a lowly resource, but if given enough attention can “turn dust into gold” and effect significant changes in the lives of many sorts.

Keywords— Animal wastes, Bio-organic fertilizer, Carabao manure tea, Dried carabao manure.

Introduction

Nutrient management and insect pest control are indispensable in any farming endeavor. Undoubtedly, the use of most appropriate technology suited to a given crop largely contributes to its growth and yield.

The utilization of indigenous farm resources such as farm manure and botanicals relative to crop management as an organic production technology has a good number of assured chain of benefits which are: environmental sanitation or ecological conservation, reduction of production cost resulting to bigger income, and production of genuinely safe food crops to address and promote health and wellness concerns.

Carabao manure is a very good bio material for enhancing soil fertility. This bio-organic resource can be collected in every farm. Rosco, 2014, (personal communication), mentioned that a carabao produces ten (10) kilos of manure a day. Chemical analysis shows that carabao manure has 18-20 C:N ratio. It has 24-30% carbon, 0.8-2.2 % Nitrogen, 2.07 Phosphorus, and 0.12- 2.0 Potassium (Aganon, et al., 2011). They further cited that manure from ruminants is found to enhance plant growth because of the decomposing microorganisms in the rumen and hormones in their urine.

With these, the study was conceived to test the efficacy of carabao manure as fertilizer as bio fertilizer on the growth and yield of cauliflower grown pots.

Methodology

Experimental Design and Treatments. A randomized complete block design (RCBD) with three replications was used. Three doses of bio-organic fertilizer from animal wastes were evaluated, to wit: No carabao manure (DCMO -control); 1.0 kg dried carabao manure (DCM1) and 1.5 kg dried carabao manure (DCM1.5). Carabao manure tea was applied as booter to all treatments. The treatments were designated as follows:

T1: DCMO - No Carabao Manure (DCMO-Control) + Carabao Manure Tea

T2: DCM1 - 1.0 kg Carabao Manure + Carabao Manure Tea

T3: DCM1.5 - 1.5 kg Carabao manure + Carabao Manure Tea

Preparation and Application of Carabao Manure as Bio-organic Fertilizer. Carabao manure was processed in two ways: dried (DCM) and manure tea (CMT). Dried carabao manure was collected and air-dried until it was totally dried, while carabao manure tea was processed by submerging a sack-full of partially dried carabao manure in a 200-liter capacity plastic drum filled with ground water for one week before its use.

Dried carabao manure (DCM) was prepared in two levels: 33 and 50 t/ha converted into 1.0 kg (DCM1) and 1.5 kg (DCM1.5), respectively. Application of DCM was done once for DCM1 and twice for DCM1.5. Application of DCM1 was administered as basal before planting; while second application for DCM1.5 as top dress was done 10 days after the first application. Carabao manure tea (CMT) at the rate of 100 ml plant was applied in each plant at planting and at weekly interval thereafter throughout the duration of the study. Application was directed in the soil at the base of each plant.

Planting and Harvesting. Hardened cauliflower seedlings (Farmers' Extra Early variety) were transplanted to the plastic pots filled with desired volume of soil medium and bio-organic fertilizer late in the afternoon to reduce transplanting shock. The plants were maintained using recommended cultural management practices. Growth was measured in terms of plant height, length and width of leaves, leaf area and weight at harvest, weight of whole plant and root-shoot-ratio. Data gathered were analyzed employing the analysis of variance (ANOVA) and treatment mean differences were determined using the Least Significance Difference (LSD) test at 5% level of probability.

Results and Discussion

Growth Characters

Growth characters as influenced by the application of carabao manure as bio-organic fertilizer are summarized in Table 1.

Plant Height at Harvest (cm). Plants applied with 1.0 kg dried carabao manure (DCM1) were tallest, 39.83 cm followed by plants applied with 1.5 kg dried carabao manure or (DCM1.5) with 38.35 cm. Plants were shortest in pots without dried carabao manure (DCM0) at 35.05 cm. Although carabao manure is considered a slow-release organic fertilizer, its effect might have been enhanced by the application of carabao manure tea (CMT), boosting plant nutrients at early crop stage resulting to the rapid growth of the cauliflowers. Jigmei, N.J., et al., (2015) reported an increase in growth parameters of broccoli applied with chicken and manure tea; and El-Magd, et al., (2016), noted accelerated growth of the plants applied with organic manure.

Number of Leaves per Plant at Harvest. Although the number of leaves produced per plant varied according to carabao manure application, the same was not significant. The production of almost the same number of leaves with or without the application of different levels of DCM indicates that the cauliflower variety used in the study can adapt to local conditions even in soils with low fertility level. This observation is true in almost all vegetable (Brassica) researches in the area where the control (without fertilizer application) plants were able to develop considerable number of leaves but of different length, breadth and weight subject to available nutrients in the soil (Tuan, 2002).

Leaf Length at Harvest (cm). Length of cauliflower leaves was significantly affected by the application of carabao manure. Longest leaves were found in DCM1 (19.55 cm) and DCM1.5 (21.02 cm); shortest leaves were noted in DCM0 (15.32 cm). Higher rates of carabao manure produced bigger and longer leaves. This implies that DCM added with CMT provided adequate nutrients to the cauliflowers. This corroborates the findings of Fayed (2010) who claimed that CMT significantly increased the vegetative parameters of the Roghini olive trees. In addition, Eltantawy (2009) also

found out that farmyard compost tea increased the height and leaf area of potato plant.

Leaf Width at Harvest (cm). Plants applied with DCM1 and DCM1.5 produced the widest leaves while those applied with DCM0 only had the narrowest leaves. The production of wider leaves in cauliflowers is necessary to support flower/curd formation. Results of this study revealed that formation of bigger and wider leaves is directly related to the application of DCM with CMT as sources of nutrient. Gross et al. (2008), reported that ammonium is the major form of nitrogen present in the extract solutions from all manure types and that the nitrogen released after 14-day extraction by the different methods from the different manures ranged between 50% and 85%. Alo and Tuan (2015) obtained almost similar result in cauliflower applied with chicken dung.

Leaf Area at Harvest (cm). The application of DCM significantly influenced the leaf area of cauliflower plants. Highest leaf area was recorded among plants applied with DCM1 with 654.26 cm² and those with DCM1.5 with 632.84 cm². Lowest leaf area was observed in the control plants (DCM0) with 403.71cm². No significant difference in leaf area was observed between DCM1 and DCM1.5. The variation in leaf area observed among treatments being bigger in plants applied with DCM than those in control plants (DCM), implies that DCM has sufficient nutrients. Nourishment received by the plants enabled the production of bigger and wider leaves; a plant requirement that allow optimum light interception and distribution for efficient physiologic process such as photosynthesis and carbohydrate transport (El-Sawy et al., 200; el-Dissoky, 2008) which contribute to the formation of reproductive parts in cauliflower.

Leaf Weight at Harvest (g). Different rates of DCM had significant influence on the weight of cauliflower leaves per plant. Average leaf weight per plant was heaviest in DCM1 at 381.39 g, and DCM1.5 with 375.55 g, but no significant difference was observed between them. Lightest leaves were observed in plants not applied with DCM0 (213.09 g). The variation in leaf weight among treatments is directly related to the size of the leaves (leaf length, width, area) produced by the plants. Based on the other growth parameters, it can be noted that the longest and widest leaves and biggest leaf area were produced by the plants treated with DCM. Gross et al., 2008, mentioned that organic fertilizers have the property to enhance soil aggregation, soil aeration, and water holding capacity; factors which offer good environmental condition to broccoli plants.

Weight of Whole Plant at harvest. Weight of whole plants differed significantly as influenced by the application of DCM. Average weight of whole plants applied with DCM1 was 235.83 g and DCM1.5 with 221.20 g were statistically the same except on plants without DCM with 139.12 g. Weight of whole plant comprises all plant parts including the roots. The results of the study relative to the weight of the whole plant almost follow

similar pattern with the other growth parameters evaluated such as the vegetative parts (leaf length, width and leaf area). The increase in vegetative growth can be traced back to the positive effects of DCM with CMT which have contributed to the increase in photosynthetic activity and uptake of soil nutrient. These results are in agreement with the findings of Caser (2009) in cauliflower, Elkhatab, (2009) on common bean.

Shoot-Root Ratio. Shoot-root ratio was highest in plants treated with DCM1.5, 8.33 g followed by plants treated with DCM1.0 with 8.85 g and those without DCM at 6.77 g per plant. Although highest shoot- ratio was highest in plants applied with DCM1.5 followed by plants applied with DCM1 and those without carabao manure (DCMO), in the descending order, the differences were not significant. The higher shoot-root ratio among treatment treated with organic fertilizer is indicative that the plants were receiving sufficient nutrients to support a large vegetative (shoot) growth. Wright et al. (1995), observed that maximum root growth and rooting depth of barley crop were higher in treatments, which received animal manures relative to where manure was not applied. The results therefore suggest that the observed response was largely due to increased availability of N and P in Carabao manure and manure tea and consequently enhanced root growth.

Table 1. Growth characters of cauliflower as influenced by the application of carabao manure.

Mean*								
Treatments	Plant Height (cm)	Number of leaves per plant	Leaf length at harvest (cm)	Leaf width at harvest (cm)	Leaf area at harvest (cm ²)	Leaf weight at harvest (g)	Weight of whole Plant (g)	Plant Root-shoot Ratio
DCMO	35.05b	16.18	15.32b	9.18b	134.57b	71.03b	19.87b	6.77
DCM1	39.83a	17.06	19.55b	11.38a	218.09a	125.18a	27.24a	7.42
DCM2	38.35a	17.06	21.02a	11.83a	210.95a	127.13a	25.58a	8.33

*Means followed by the same letter are not significant at 5% DMRT.

Conclusion and Recommendations

The application of carabao manure complemented with carabao manure tea as source of bio-organic fertilizer for cauliflower production was found to have highly influenced the growth of cauliflowers, evident on the different growth parameters evaluated, such as plant height, length and width of leaves, leaf area and weight at harvest, weight of whole plant and root-shoot-ratio. Carabao manure with manure tea was found to be effective in influencing the growth of cauliflower. Higher rates of carabao manure enhanced with manure tea resulted to taller, bigger and heavier cauliflower plants.

Carabao manure and carabao manure tea, a farm waste, is rich in plant nutrients and beneficial microorganisms are a potential bio-organic fertilizer productive and sustainable organic cauliflower production. A follow-up study is necessary to further verify the performance of cauliflower applied with carabao manure with manure tea under farmers field conditions.

Acknowledgement

The researcher wishes to express her profound thanks to the University of Eastern Philippines (UEP) administration and the Commission on Higher Education (CHED) for the material and financial support that leads in the realization of this experiment. To the University Research and Development Services staff, and field workers who assisted the researchers throughout the duration of the experimentation.

Bibliography

1. Abou El- Magd, M.M., A.M, El-Bassiony and Z.F. Fawzy. 2016. Effect of Organic Manure with or Without Chemical Fertilizers on Growth, Yield and Quality of Some Varieties of Broccoli Plants.
2. Aganon, Clarita P., Nestor D. Pagaduan, Luzviminda M. Galang, Tomas D. Gajete, Rex S. Parayno, Arturo M. Alegado, Rafael R. Rafael, Fe L. Potenciuncula, Matilde Mellicent S. Recto, Pablo J. Rafael, Jr. and Gerry Z. Laggui. 2011. Biofertilizer Production and Entrepreneurship Training Manual. Central Luzon State University. Science City of Munoz, Nueva Ecija: CLARRDEC, CVARRD and PCARRD-DOST. 92p.
3. Alo, L.D. and Tuan, P.P. 2014. Growth and Yield of Cauliflower (*Brassica oleracea*) Applied with Chicken Dung-based Organic Fertilizer under Polyethylene Pot Culture. UEPGS MS Thesis.
4. Caser G. Abdel.2009. Improving the Production of Well Irrigated Cauliflower (*Brassica oleracea* var. Botrytis, Cv. Snowball Y. Imp) by Foliar Spraying of some Growth Regulators. Journal of ZankoySulaimani12(1), 29-49.
5. DOST. (1998): Madre de cacao. Department of Science and Trade. Manila, Philippines. Stoll, G. (2000): Natural protection in the tropics. MargrafVer Weikersheim.www.bar.gov.ph
6. El-Dissoky, R. A. 2008.Studies on the use efficiency of potassium and organic fertilizers on potatoes and its role in improving soil properties. PhD Thesis. Faculty of Agriculture, Mansoura, University, Egypt.
7. Elkhatab HA. 2009. Growth and yield of common bean (*Phaseolus vulgaris* L.) in response to Rhizobium Inoculation, nitrogen and molybdenum fertilization. Alex. Sci. Exchange Journal 30, 319- 332.
8. El-Sawy, B. I., Radawan, E. A. and Hassan, N. A. 2000. Growth and yield of potato as affected by soil and foliar potassium application. Journal of Agriculture Science, 25(9):5843-5850.
9. El-Tantawy, I. M., El-Ghamry, A. M. and Habib, A. H. 2009. Impact of farmyard manure and manure compost tea on potato yield and soil fertility. Journal of Agriculture Science, 34(1): 669 – 678.
10. Fayed, T. A. 2010. Optimizing yield, fruit quality and nutrient status of Roghiani olive grown in Libya using some organic extracts. Journal of Horticulture Science and Ornamental Plants, 2(2): 63-78.

11. Gross, A., Arusi, R. and Nejidat, A. 2007. Assessment of extraction methods with fowl manure for the production of liquid organic fertilizers. *Bioresource Technology*, 99: 327– 334.
12. Jigme¹, Nipon Jayamangkala¹, Pathipan Sutigoolabud, JiraponInthasan and Siriwat Sakhonwasee¹. 2015. The effect of Organic fertilizers on growth and yield of Broccoli (*Brassica oleracea* L. var. *italica* Plenck cv. Top Green) *Journal of Organic Systems*, 10(1),
13. PCARRD. 1999. Broccoli and Cauliflower Production Guide. Information Bulletin No. 148. Los Banos 4030 Laguna.
14. Rodriguez AA. January 2012. "Sariaya, Quezon Organic Farmer Succeeds with Cauliflower". *Agriculture Monthly*. Volume 16, Issue 01. p. 44.
15. Samonte P. August 2012. "Bataan Farmer earns Millions from Calamansi and Vegetables". *Agriculture Monthly*. Volume 16, Issue 8. pp.20-21[15]
Samonte P. March 2013. "Organic Farming Develops Roots In Pampanga". *Agriculture Monthly*. Vol 17, Issue No. 3 pp 52-53.